




15th INTERNATIONAL SEDIMENTOLOGICAL CONGRESS



INTERNATIONAL
ASSOCIATION OF
SEDIMENTOLOGISTS



Universitat d'Alacant
Universidad de Alicante



15th INTERNATIONAL SEDIMENTOLOGICAL CONGRESS



APRIL 12-17, 1998

Abstracts

Editors:

Juan C. Cañaveras
M. Ángeles García del Cura
Jesús Soria

SCIENTIFIC COMMITTEE

Gaspar Alonso Gavilán	Univ. Salamanca, Spain.	Javier Martín Chivelet	Univ. Complutense de Madrid, Spain.
Ana M. Alonso Zarza	Univ. Complutense de Madrid, Spain.	Peter I. Martini	Univ. Guelph, Canada.
Pere Anadón	Inst. Jaume Almera, CSIC, Spain.	Mariana Marzo	Univ. Barcelona, Spain.
Alfreda Arche	Inst. Geol Económica, CSIC, Spain.	Judith A. McKenzie	Geol. Inst. Zürich, Switzerland.
Alwyn E. Annels	Univ. Kordiff, UK.	José M. Molina	Univ. Jaén, Spain.
Julian E. Andrews	Univ. of East Anglia, UK.	J.A. Muñoz	Univ. Barcelona, Spain.
José Arribas	Univ. Complutense Madrid, Spain.	Philippe Muchez	Univ. Leuven, Belgium.
Victor R. Baker	Univ. of Arizona, USA.	Emiliano Mutti	Univ. Parma, Italy.
Teresa Bardaji	Univ. de Alcalá de Henares, Spain.	L.M. Nieto	Univ. Jaén, Spain.
Thilo Bechstaedt	Univ. Heidelberg, Germany.	Gary Nichals	Univ. London, UK.
Danw. J. Bosence	Royal Holloway Univ. of London, UK.	Federio Ortí	Univ. Barcelona, Spain.
Juan C. Braga	Univ. Granada, Spain.	Cecile Pabian Goyheneche	Elf Explor. Prod. Pau., France.
Luis A. Buatois	Kansas Geol. Survey, USA.	Chris Paola	Univ. Minnesota, USA.
Manuel Bustillo	Univ. Complutense Madrid, Spain.	Albert Permyer	Univ. Barcelona, Spain.
David A. Cacchione	U.S. Geological Survey, USA.	Tadeusz M. Peryt	Univ. Warsaw, Poland.
Juan C. Cañaveras	Univ. Alicante, Spain.	A. Guy Plint	Univ. Western Ontario, Canada.
Henry S. Chafetz	Univ. of Houston, USA.	Lluis Pomar	Univ. Islas Baleares, Spain.
Maria Bianca Cita	Univ. Milana, Italy.	George Postma	Univ. Utrecht, The Netherlands.
Lluis Cabrera	Univ. Barcelona, Spain.	Cay Puigdefábregas	Nork Hydro Res. Centre, Norway.
Lars B. Clemmensen	Univ. of Copenhagen, Denmark.	Victoriano Pujalte	Univ. País Vasco, Spain.
Jasé R. Colmenero	Univ. Salamanca, Spain.	Harold G. Reading	Univ. Oxford, UK.
Ferran Colomba	Univ. Barcelona, Spain.	Robert Riding	Univ. Kordiff, UK.
Carmen Comas	Univ. Granada, Spain.	Franciso J. Rodríguez Tovar	Univ. Granada, Spain.
Marinos E. Danselaar	Univ. of Delft., Netherlands.	Jean Marie Rauchy	CNRS, France.
Javier Elorza	Univ. País Vasco, Spain.	David Rubin	U. S. Geological Survey, USA.
Mateu Esteban	Univ. Islas Baleares, Spain.	Pedro A. Ruiz Ortiz	Univ. Jaén, Spain.
Chris Fielding	Univ. Queensland, Australia.	Carlos Santiesteban	Univ. Literaria de Valencia, Spain.
Gerald M. Friedman	Northeastern Sci. Foundation, USA.	Walther Schwarzacher	The Queen's Univ. of Belfast, UK.
Peter F. Friend	Univ. Cambridge, UK.	Jordi Serra	Univ. Barcelona, Spain.
Vicente Gabaldón	ITGME, Spain.	Tsunemasa Shiki	Univ. Kyoto, Japan.
M.A. García del Cura	Inst. Geol. Económica, CSIC, Spain.	Alfonso Sopena	Inst. Geol. Económica, CSIC, Spain.
José C. Garcia Ramos	Univ. Oviedo, Spain.	Jesús M. Soria	Univ. Alicante, Spain.
Guillermina Garzón	Univ. Complutense de Madrid, Spain.	Michael G. Stamatakis	Univ. Athens, Greece.
Robert H. Goldstein	Univ. Kansas, USA.	Daniel J. Stanley	N. Mus. Nat. History Washington, USA.
Janos Haas	Univ. Budapest, Hungary.	Cantxita Taberner	Inst. Jaume Almera, CSIC, Spain.
Bilal Haq	Nat. Science Found. Arlington, USA.	Michael G. Talbot	Univ. of Bergen, Norway.
Adrian Harvey	Univ. Liverpool, UK.	Torbjorn E. Törnqvist	Univ. Utrecht, Netherlands.
Ian Jarvis	Kingston University, UK.	Trinidad de Torres	Univ. Politécnica de Madrid, Spain.
Noel P. James	Queen's Univ. Ontario, Canada.	Maurice E. Tucker	Univ. Durham, UK.
Brian G. Jones	Univ. Wollongong, Australia.	Chris Vasconcelos	Geol. Inst. Zürich, Switzerland.
Wolfgang E. Krumbein	Univ. Oldenburg, Germany.	Mario Vendrell	Univ. Barcelona, Spain.
Antonio C. Lasaga	Univ. Yale, USA.	Jaume Vergés	Univ. Barcelona, Spain.
Bernard Long	Univ. Quebec, Canada.	Lorenza Vilas	Univ. Complutense de Madrid, Spain.
Romón Mas	Univ. Complutense de Madrid, Spain.	Federico Vilas	Univ. Vigo, Spain.
Eudald Maestro	Univ. Autónoma de Barcelona, Spain.	Roberto H. Wagner	Jardín Botánico de Córdoba, Spain.
Jose M. Martín	Univ. Granada, Spain.	V. Paul Wright	Univ. Kordiff, UK.
Carmen Moreno	Univ. Huelva, Spain.	Gian Gaspare Zuffo	Univ. Bologna, Italy.

SCIENTIFIC PROGRAMME ACCORDING TO SESSIONS

THEMES

THEMES A. ANALYTIC TECHNIQUES. DATA PROCESSING AND REPRESENTATION

J. Arribas, J.A. de la Peña & A. Tortosa: Detrital modes in sedimentoclastic sands from first-order streams of the Iberian Range, Spain: the effect of source lithology	153
S.M. Bidart: Artificial laboratory aggregates in sediments	197
M. Canals, B. Alonso, J. Sorribas, M. Farran & Big '95 Team: Coupled swath bathymetry and ultra-high resolution seismic parametric profiling reveal sedimentary processes and allow precise volume calculations of channel-levee complexes.....	227
M.C. Castellano, G. Nardi, M. Piscitello & M.L. Putignano: Anisotropy of magnetic susceptibility and sedimentological analysis. a multidisciplinary approach to the study of a synorogenic basin	241
R. Chandrajith, H.J. Tobschall & C.B. Dissanayake: Distribution of REEs and selected trace elements in grain size fractions of sediments from Walawe Ganga River, Sri Lanka	250
S. Fernández-Bastero, L. Gago-Duport, O. Pazos, I. Alejo, B. Rubio, D. Rey, S. García-Gil, F. Vilas & A. Santas: Chemical and structural markers of the glaucony maturity. Results from syntheses experiments and XRD Rietveld analysis	328
A.D. Kirkbride, J.L. Best, T. Buffin-Belanger & I. Reid: Monitoring, processing and presenting space-time turbulence data: new opportunities from ultrasonic velocity profiling	473
H. Krawinkel, S.Wozazek, J. Krawinkel & W. Hellmonn: Heavy-mineral analysis and clinopyroxene geochemistry applied to provenance analysis of lithic sandstones from the Azuero-Soná Complex (NW Panamá)	478
S. Montesinos & J. Arribas: Source area versus detrital products: a geographical information system approach	558
C. Reiser, A. Lejay & G. Dromart: Automatic recognition of sedimentary bodies and genetic sequences by analysis of wireline log shape.....	654
D. Scradeanu, C. Petrache & D. Jipa: Geostatistical analysis of Upper Neogene lacustrine deposits of the Dacian Basin, Romania	708
D. Sherman & E. Farrell: The Charnock relationship for aeolian saltation	719
M. Sirat, A. Aldahan & U. Nordlund: A neural networks approach to predict sedimentation and climate changes in arctic sediment covering last 350ka	725
T. Slomka & E. Slomka: Modal and model sequences in Carpathians flysch sedimentary series using Markov chains analysis	727
D.C. Sousa & H.Vital: Stratigraphic significance of heavy minerals of the sediments from NE Brazil	733
K.-W. Tietze, D. Bannert, F. Böker, F. Diederich, B. Dietrich, M. Karle, F. Lemmer, D. Radies & E. Rose: Macrotidal bedforms as seen in the thermal infra-red	763
M.C. Zuluaga, F. Carcía-Garmilla & J. Arostegui: Different paths of provenance for the Middle Albian Gordexola Formation sediments (Basque-Cantabrian Basin, northern Spain)	851

R. Salas, J. Guimerà, J. Giménez-Montsant & C. Martín-Closas: Mesozoic rift structure, sedimentation and Palaeogene inversion of the intraplate extensional Maestrat Basin. Iberian Range (NE Spain)	689
Y. Sánchez-Moya & A. Sopena: Tectonic sequences and fluvial architecture. The Mesozoic onset infill of an extensional basin (Central Spain)	694
A. Schäfer, R.J. Korsch & R. Müller: Permo-Carboniferous Soar-Nahe Basin (Germany), structure and molasse basin-fill ..	702
Liu Shanyin & Zhong Daloi: Tectono-sedimentary development of Tertiary Basin in Jinggu-Zhenyuan region, SW Yunnan, China	717
A. Siedlecka, V.G. Olovyanishnikov & D. Roberts: Neoproterozoic passive-margin sedimentation along the NE Perimeter of the east European Craton	721
S. Sierra & C. Moreno: Alluvial fans as markers of Viar basin early evolution (SW Spain)	722
S. Sierra, C. Moreno & M.A. Casalsvázquez: Petrographic evidences of tecto-sedimentary activity in the Viar Basin (SW Spain)	723
K. Strand: Sedimentary facies and sediment composition changes in a rift/transform margin transition	743
O. Takano: The changes of depositional systems and sequences in response to basin evolution in a rifted and inverted basin, Central Japan	752
J.H. Ten Veen & G. Postma: Reconstruction of fault block kinematics and architecture of neogene outer-arc basins of central and eastern Crete (Greece) -integration of satellite, topographic and field data into a GIS	758
J.H. Ten Veen, G. Postma, H. de Boorder, F. Slangen, A. Werre, S. Van Ratingen & V. Unnithan: Reconstruction of fault block kinematics by integration of land-sat, topographic and stratigraphic data into a GIS	759
C. Tiratsoo: Basement uplift and sedimentary basin development in the eastern Betic Cordillera, SE Spain	764
A. Tortosa & J. Arribas: Evolution of sandstone composition in a continental foreland basin. Loranca Basin, Spain	769
M. Tropeano, L. Sabato, P. Pieri & C. Doglioni: Geodynamic and stratigraphic history of the Bradanic Trough (South Apennines Foredeep - Italy)	772
B.R. Turner & K. Thompson: Triassic tectono-stratigraphic events in the main Karoo Basin, South Africa, and their relation to the Falkland Plateau: a new model for provenance tectonic activity	777
D. Uličný: Interplay of strike-slip tectonics and eustasy in shallow-marine clastic wedges, Bohemian Cretaceous Basin ..	778
A. Valente: Tectono-sedimentary evolution of the Cilento Basin (Italy)	780
A. Voznesensky, A. Knipper, A. Perfiliev, E. Uspenskaya & A. Areshin: Middle to Late Jurassic history of the eastern part of the Mountain Crimean terrain	808
M. Wendarff: Olistostromes in the Lufilian Arc, Central Africa	815
U. Wortmann & H. Weissert: Climate & plate-tectonics: Climate & plate-tectonics: Early Cretaceous sandstone sequences of the western Tethys	825
L. Zitouni, M. Bédim & S. Tlig: Holokinesis and subsurface geology of Mesozoic basins in central Tunisia	845

THEME G. GEOLOGY AND OCEANOGRAPHY OF CONTINENTAL MARGINS

J. Acosta, P. Herranz, A. Muñoz, C. Palomo, M. Pardo de Domlebum & E. Uchupi: Channel geomorphology of Columbrete Island area. Valencia Margin	114
--	-----

formations seems to be the more sensitive sedimentary Lithology to the gradient value, increasing its SGI when this value increases. Finally, the effect of weathering on siliciclastic material can be evaluated comparing the K/Q+K index in sandstone formations (0.27 - 0.19) and in the sands (0.11 - 0.08), representing a decrease in feldspar of 15% aprox. with regard to total siliciclastics grains of the sediment.

COEVAL CARBONATE GRAINS IN MODERN FLUVIAL SANDS (IBERIAN RANGE, SPAIN): THE ACTIVITY OF A CONTINENTAL CARBONATE FACTORY

M^a. E. Arribas & A. Tortosa

Departamento de Petrología y Geoquímica. Universidad Complutense. 28040-Madrid, Spain

Petrographic analysis in head stream sands derived from Mesozoic siliciclastic and carbonate rocks in the Iberian Range show a variable content in coeval carbonate material. These components are bioclasts, micritic and sparitic grains and coatings around grains. Generally, their origin is related to the erosion of recent carbonate deposits as cool freshwater tubas, calcimorphic paleasols and speleothems. These carbonate deposits can be mainly associated to carbonate precipitation in a karstic environment developed in several formations outcropping in the work area. This study is a tentative characterisation of this coeval carbonate material based on a detailed petrographic analysis in coarse, medium and fine sand fractions.

The presence of coeval carbonate grains in the sediments produces an increase in volume of the finer sand fraction, where these components tend to concentrate. Occasionally their content constitute more than 90% of sediment volume in this fraction. Whereas siliciclastic and carbonate grains derived from the parent rocks prevail in the coarse sand fraction. These grains show abundant coatings of coeval carbonate with variable textures.

Textural and compositional criteria permit to describe four categories of coeval carbonate grains: micritic, sparitic, coated and bioclasts. The most frequent coevals are micritic grains which show a spongy and porous microstructure, sometimes preserving isolated algal filaments or algal colonies of fan-like character. The origin of these coeval grains are associated to the erosion of recent tufa deposits. Also laminated micritic grains occur as ancolite (coated grains) or stromatolite remains. Other kind of micritic coeval are paleosol fragments which are recognised as hard micritic grains. The sparite grains are abundant and generally are the result of algal micrite sparitization. These grains can preserve a fan-like microstructure formed by algal filaments (similar to those described in micritic grains) or a prismatic and radial microstructure that characterises *Microcodium* colonies. In addition, sparite grains without microstructure appear like isolated or cluster crystals with an uncertain origin. The coated grains are very common and are constituted by a nucleus plus a coating of coeval carbonate (micritic and/or sparitic coatings). When the nucleus correspond to extrabasinal grains, as quartz, feldspar or carbonate rock fragments, this kind of grain had been named as compound grains (Arribas *et al.*, in this meeting), and when the nucleus are intrabasinal in origin we have named as coated coeval grains. Coatings are formed by micritic or sparitic carbonate in as single or multisteady envelopments and their origin is related with algal activity that may form algal-mots on the channelized deposits. The identified bioclasts are charophytes (stems and small reproductive gyrogonites) and gastropods.

Although these coeval components have been well defined in basis on textural and compositional criteria, some problems exist when grains with different origin show similar microfabrics. This convergence of microfabrics is common in grains with a laminated microstructure as: debris of stromatolites, oncolites, rhizocretions and speleothems. In addition, early diagenetic processes as cementation or recrystallization, are very frequent in continental environments. These processes may produce an increase of crystal size of carbonate deposits, and then may provide coeval grains with very similar textures to those grains derived from carbonate source rocks. This problem is greatly dramatised in petrographic analysis carried out on ancient sandstones deposits, when general diagenesis mask the primary textures. Classically, authors have used textural and compositional criteria as: coeval bioclasts, soft appearance, irregular contours, grain-size coarser than the detrital grains and lack of recrystallization and cementation processes to identify intrabasinal grains in ancient deposits. These studies underestimate the presence of coeval carbonate grains affected by early recrystallization and cementation. Probably these grains were considered as false extrabasinal components.

Also in the fine fraction an important accumulation in micritic grains without microstructure occur as consequence of the loss of textural characters when grain size decreases. For these reasons is advisable the analysis of textures of coeval carbonate material in coarse fractions.